HERBICIDE GUIDE



Active Ingredients in Herbicides - sometimes formulated in combination

TYPE	USE/TARGET	
Glyphosate	Roundup, Gligel, (cut'n'paste) Glyphosate 360/510, non selective	
Metsufuron- Metyl	Escort, Mustang, Meturon, Metgel (cut'n'paste), non selective	
Picloram	Picloram gel (cut'n'paste), non selective	
Haloxyfop	Gallant, Haloxyfop 100, selective, targeting grasses	
Triclopyr	Grazon, Brushoff, Triclop 600, non selective	
2-4D	BanVine/ LawnGuard, Yates Woody Weed Killer, selective broad leaf	
Amitrole	Activated Amitrole, non selective	
Dicamba	Oracle, LawnClean, selective broad leaf weed control	

There are also penetrants or crop oils to make spray stick better and dye products to mark sprayed areas Toxicity class on a label:

- Class 6 relates to vertebrates,
- 6A being the most toxic
- Class 9 means Ecotoxic, affecting water ways





Gligel or glyphosate in a sprayer for:

- Gorse,
- Tabacco weed (drilling best),
- Climbing asparagus,
- Pampas

Metsulfuron:

- Cotoneaster,
- Blackberry,
- Mignonette vine,
- Moth Plant,
- Japanese Honey suckle

IS THE PLANT FRUITING

Are you trying to hit a vine such as moth plant or passion fruit that has already got fruit on? Try a special mixture to paint on the bottom 50cm of the stems to lower the germination of seeds:

- Mix 250ml of water, 0.25 teaspoons/ 1gr of Metsulfuron powder, add 20ml of cooking oil (any) in a screw top jar and shake well.
- Use a narrow paint brush to paint on, making sure oil is not separating on top again before applying.

WHICH HERBICIDE FOR WHICH INVASIVE WEED

Invasive plant	Herbicide options	Ratios	Ideal time of year
Wooly Nightshade	Glimax on cut		Any time of year but remove seeds
	Or Glyphosate	Undiluted for drill & fill	beforehand
Wattles	Triclopyr	Undiluted for drill & fill	Late winter or Spring,
	or Metgel	For stump painting	when flowering
Taiwan cherry	Glimax	Undiluted for drill& fill	Late winter or Spring, when flowering
	Or Metsulfuron- Methyl	1g per litre	
Wildling conifers	Glyphosate	Undiluted, Drill & frill	Any time of year but late winter or Spring
	Or Metsulfuron- Methyl	20g per litre	before cones develop
	Or Xtree basal	Undiluted, paint on	
		bark	
Elaeagnus	Metgel	For stump painting	Any time of year but preferably before
	Or Picloram	For stump painting	seeds set
	Brush killer		
Privets	Metgel	For stump painting	Any time of year but preferably before
			flowering/ seeds set
Gorse	Glimax	For stump painting	Any time of year but avoid spreading
	Or Metgel	For stump painting	seeds
Cotoneaster	Metgel	For stump painting	Before seeds set in Autmn
	No alternative		
Cestrums, eg Queen	Metgel	For stump painting	Winter when trees start flowering
of the night	Or Triclopyr	15ml per litre	
Blackberry	Metgel	For stump painting	Before berries set
	Triclopyr		

Passion vines	Glyphosate	For stump painting	Summer before fruit develops
	Or Metgel		
Mignonette vine	Metsulfuron- Methyl	5g/ 10 litres of water	Summer before fruiting bodies form
	Or Triclop		
Japanese honey	Metgel	For stump painting	Spring when it flowers
suckle	Gligel less effective		
Moth plant	Metgel	For stump painting	Any time of year but preferably before
	Or Metsulfuron- Methyl	5g/ 10 litres of water	pods develops late Summer
	Brushkiller or Banvine		
Blue morning glory	Glyphosate	100ml/ 10 l of water	Flowering time when visible
	Or Glimax	For stump painting	
	Or Metgel	For stump painting	
Jasmine	Metsulfuron- Methyl	5g/ 10 litres of water	Flowering time when visible
	Or Glyphosate	100ml/ 10l of water	
Climbing Aspargus	Glyphosate	3.3- 10ml per litre, no	Any time of year but poss. remove seeds
		penetrant to protect	beforehand
		native plants	
Wild Ginger and most	Metsulfuron- Methyl	2g for 10 litres of water	Any time of year but easier when it
other bulbs eg Agap.	No alternative		flowers

For more information contact us on: WEEDACTION@PIROACONSERVATION.ORG.NZ

